

# Antibacterial Activities of Iron Chelators against Common Nosocomial Pathogens

Mitchell G. Thompson, Brendan W. Corey, Yuanzheng Si, David W. Craft,\* and Daniel V. Zurawski

Walter Reed Army Institute of Research, Silver Spring, Maryland, USA

The activities of iron chelators (deferoxamine, deferiprone, Apo6619, and VK28) were evaluated against type strains of *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, and *Escherichia coli*. Deferiprone, Apo6619, and VK28 each inhibited growth in standard and RPMI tissue culture medium, while deferoxamine had no effect. Additionally, time-kill assays revealed that VK28 had a bacteriostatic effect against *S. aureus*. Therefore, these newly developed iron chelators might provide a nontraditional approach for treatment of bacterial infections.

Iron is an essential cofactor of biochemical pathways in both prokaryotic and eukaryotic species. Numerous studies have assessed the potential viability of iron chelators as therapeutic agents against various microbes, but with only mixed success (4, 6–8, 10–17, 20, 24, 25, 27). Nevertheless, as novel iron chelators are developed for treatment applications such as neurodegenerative diseases (3, 9, 19) or  $\beta$ -thalassemia (3, 9), an evaluation of their antimicrobial activities should be tested, because their efficacies against bacteria may be superior to chelators previously tested. In the case of multidrug-resistant (MDR) bacteria, where entire classes of antibiotics are no longer treatment options (1, 21), iron chelators that have already undergone toxicity and preclinical testing in animals might provide an alternative treatment approach. MDR species, such as *Staphylococcus aureus* or *Acinetobacter baumannii*, are exceedingly difficult to treat because of nosocomial spread and infections in immunocompromised patients (1, 18, 21). The same microbes have also been responsible for wound infections incurred by military personnel who are immunocompromised after polytrauma (2, 23).

Defersirox and deferoxamine are approved by the U.S. Food and Drug Administration (FDA) but have demonstrated limited efficacies in combating bacterial infections (4, 6, 17). Deferoxamine is a siderophore, a molecule secreted by bacteria to capture iron; therefore, many bacteria challenged with deferoxamine also harbor a receptor capable of capturing such molecules when complexed to iron (4, 6, 17). Defersirox, while rationally designed to bind iron, failed to treat fungal infections (20) and is considered toxic (11).

(Some of this work was first presented at the 51st Interscience Conference on Antimicrobial Agents and Chemotherapy, 17 to 20 September 2011, Chicago, IL.)

In this study, we sought to assess the antibacterial effects of iron chelators that have yet to be tested against bacteria, as well as deferiprone, a chelator recently FDA approved for iron overload due to blood transfusions in patients with thalassemia. Deferiprone has also been shown to have antibacterial properties against certain bacterial species *in vitro* (4, 10). Deferiprone (ApoL1) and Apo6619 were provided by ApoPharma, Inc., and VAR10100 (VK28 dihydrochloride) was provided by Varinel, Inc. Because of their iron chelation properties, both Apo6619 and VK28 (and their derivatives) are currently being studied for treatment applications (9, 19, 22, 28). Deferoxamine mesylate salt (DFO) and 2,2'-bipyridyl (DIP) were purchased from Sigma-Aldrich Inc. and were evaluated for comparison purposes.

TABLE 1 MICs of iron chelators against ATCC type strains grown in CAMHB

Bacterial species	Strain no.	MIC of iron chelator ( $\mu\text{g/ml}$ )				
		DIP	DFO	ApoL1	Apo6619	VK28
<i>A. baumannii</i>	17978	64	>512	128	256	128
<i>A. baumannii</i>	19606	64	>512	128	256	128
<i>S. aureus</i>	25923	512	>512	>512	>512	256
<i>S. aureus</i>	43300	256	>512	>512	>512	256
<i>P. aeruginosa</i>	PAO1	256	>512	256	>512	>512
<i>P. aeruginosa</i>	27853	256	>512	>512	>512	>512
<i>K. pneumoniae</i>	BAA-2146	256	>512	256	>512	>512
<i>K. pneumoniae</i>	700603	512	>512	512	>512	>512
<i>E. coli</i>	35718	64	>512	512	256	>512
<i>E. coli</i>	43888	64	>512	512	256	>512

Bacterial type strains considered common nosocomial infectious agents were acquired from the American Type Culture Collection (ATCC): *A. baumannii* (19606 and 17978), *Pseudomonas aeruginosa* (PAO1 and 27853), *S. aureus* (43300 and 25923), *Klebsiella pneumoniae* (BAA-2146 and 700603), and *Escherichia coli* (35718 and 43888). The MICs of DIP, DFO, ApoL1, Apo6619, and VK28 were determined by following the microdilution methodology recommended by the CLSI (5) in cation-adjusted Mueller-Hinton broth (CAMHB) against the bacteria listed above. The MICs were also determined in RPMI 1640 tissue culture medium (Life Technologies, Inc.), which may better represent the human host environment, with limited amounts of cofactors such as calcium, magnesium, zinc, and iron. Time-kill assays against *S. aureus* and *E. coli* were performed as described by White et al. (26). An initial inoculum of  $\sim 1.0 \times 10^7$  CFU/ml was challenged with either  $1 \times$  or  $2 \times$  the MIC of VK28, and cells were grown at  $37^\circ\text{C}$  for 24 h. Samples were

Received 6 June 2012 Returned for modification 10 July 2012

Accepted 25 July 2012

Published ahead of print 30 July 2012

Address correspondence to Daniel V. Zurawski, daniel.zurawski@amedd.army.mil.

\* Present address: David W. Craft, Microbiology, Penn State Milton S. Hershey Medical Center, Hershey, Pennsylvania, USA.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.01197-12

The authors have paid a fee to allow immediate free access to this article.




FIG 1 Time-kill studies of VK28 against *Staphylococcus aureus* 43300 (A) and *Escherichia coli* 48333 (B). The no-treatment results represent growth in the absence of chelator. The detection limit was  $1.0 \times 10^3$  CFU/ml.

taken at 0, 2, 6, and 24 h, and the CFU/ml was determined via dilution and plating with a spiral plater (Advanced Instruments, Inc.), with each sample diluted over 3 logs onto plates containing CAMHB medium and agar. Biological replicates for all tests were performed at least three times in triplicate (technical replicates).

DFO did not affect bacterial growth in CAMHB (MIC,  $>512$   $\mu\text{g/ml}$  for all bacteria tested) (Table 1). The result was not surprising, since the compound may readily deliver iron to bacteria with a cognate siderophore receptor. In contrast, VK28 inhibited the growth of *A. baumannii*, *E. coli*, and *S. aureus* in CAMHB. Further, both ApoL1 and Apo6619 inhibited the growth of some strains of *P. aeruginosa* and *K. pneumoniae*, as well as *E. coli* and *A. bauman-*

*nii*, while no effect on *S. aureus* was observed (Table 1). Because CAMHB is a rich broth with excess iron, carbon sources, and other cofactors far exceeding the levels in the human body, RPMI medium was chosen to evaluate the activity on the same bacterial strains in a more restrictive medium. When challenged with the iron chelators that showed the most promise in CAMHB, the MICs were reduced accordingly in this cofactor-limited environment (Table 2). When evaluated in RPMI, for VK28 the MIC improved 4- to 64-fold, and the ApoL1 and Apo6619 MICs improved 2- to 4-fold (Table 2).

Previous studies with iron chelators have demonstrated a bacteriostatic effect on bacterial growth (10, 14, 16). We performed a

**TABLE 2** MICs of iron chelators against ATCC type strains grown in RPMI 1640 medium

Bacterial species	Strain no.	MIC of iron chelator ( $\mu\text{g/ml}$ )				
		DIP	DFO	ApoL1	Apo6619	VK28
<i>A. baumannii</i>	17978	32	>512	64	64	32
<i>A. baumannii</i>	19606	32	>512	128	128	8
<i>S. aureus</i>	25923	NA <sup>a</sup>	NA	NA	NA	32
<i>S. aureus</i>	43300	NA	NA	NA	NA	16
<i>P. aeruginosa</i>	PAO1	256	>512	128	512	16
<i>P. aeruginosa</i>	27853	256	>512	512	>512	16
<i>K. pneumoniae</i>	BAA-2146	128	>512	256	512	16
<i>K. pneumoniae</i>	700603	256	>512	256	512	16
<i>E. coli</i>	35718	64	>512	512	256	32
<i>E. coli</i>	43888	64	>512	256	256	8

<sup>a</sup> NA, not attempted.

time-kill assay to see if this was also true of the iron chelators evaluated in this study. *E. coli* and *S. aureus* were both challenged with either 1 $\times$  or 2 $\times$  the determined MIC for VK28 (Fig. 1). Growth of *S. aureus* was attenuated somewhat by both concentrations (Fig. 1A). Additionally, VK28 proved to have a bacteriostatic effect on *E. coli* at 2 $\times$  the MIC (Fig. 1B). Similar bacteriostatic effects were observed for ApoL1 and Apo6619 (data not shown).

Unlike the mild effects observed with DFO and other iron chelators in previous studies (4, 9), VK28, ApoL1, and Apo6619 had pronounced effects on the growth of nosocomial bacteria. The outcomes observed could be related to the rational design of these iron chelators. For example, VK28 includes a piperazine ring that enhances polarity to cross the blood-brain barrier for the treatment of neurodegenerative disease (19, 28). It is possible that this polarity may also allow the chelator to cross certain bacterial membranes. ApoL1, in contrast, is a very small, neutral molecule, and these properties are known to facilitate its passage across host cell membranes and perhaps bacterial membranes as well. Therefore, in each case, the free iron might be chelated both inside and outside the bacteria, explaining the enhanced efficacy. Continued studies on these molecules, including combinatorial therapies with conventional antibiotics and animal modeling, will attempt to uncover the mechanisms by which these iron chelators provide a potent antimicrobial effect.

## ACKNOWLEDGMENTS

We thank Michael Spino and John Connelly from ApoPharma, Inc., and Vincent R. Zurawski, Jr., from Varinel, Inc., for helpful discussions and critical reading of the manuscript.

The findings and opinions expressed herein belong to the authors and do not necessarily reflect the official views of the WRAIR, the U.S. Army, or the Department of Defense.

## REFERENCES

- Boucher HW, et al. 2009. Bad bugs, no drugs: no ESKAPE! An update from the Infectious Diseases Society of America. *Clin. Infect. Dis.* 48:1–12.
- Brown TS, Hawksworth JS, Sheppard FR, Tadaki DK, Elster E. 2011. Inflammatory response is associated with critical colonization in combat wounds. *Surg. Infect.* 12:351–357.
- Chakrabarty P, et al. 2011. Iron chelation therapy and thalassemia: an overview. *Mymensingh Med. J.* 20:513–519.
- Chan GC, Chan S, Ho PL, Ha SY. 2009. Effects of chelators (deferrioxamine, deferiprone and deferasirox) on the growth of *Klebsiella pneumoniae* and *Aeromonas hydrophila* isolated from transfusion-dependent thalassemia patients. *Hemoglobin* 33:352–360.
- CLSI. 2009. Performance standards for antimicrobial susceptibility testing; 19th informational supplement, p 152. Clinical and Laboratory Standards Institute, Wayne, PA.
- Collins HL, Kaufmann SH, Schaible UE. 2002. Iron chelation via deferrioxamine exacerbates experimental salmonellosis via inhibition of the nicotinamide adenine dinucleotide phosphate oxidase-dependent respiratory burst. *J. Immunol.* 168:3458–3463.
- Debebe Z, et al. 2011. Iron chelators of the di-2-pyridylketone thiosemicarbazone and 2-benzoylpyridine thiosemicarbazone series inhibit HIV-1 transcription: identification of novel cellular targets—iron, cyclin-dependent kinase (CDK) 2, and CDK9. *Mol. Pharmacol.* 79:185–196.
- Fernandes SS, et al. 2010. Identification of a new hexadentate iron chelator capable of restricting the intramacrophagic growth of *Mycobacterium avium*. *Microbes Infect.* 12:287–294.
- Heli H, Mirtorabi S, Karimian K. 2011. Advances in iron chelation: an update. *Expert Opin. Ther. Pat.* 21:819–856.
- Kim CM, Shin SH. 2009. Effect of iron-chelator deferiprone on the *in vitro* growth of staphylococci. *J. Korean Med. Sci.* 24:289–295.
- Kontoghiorghe GJ. 2011. A record of 1320 suspect, deferasirox-related, patient deaths reported in 2009: insufficient toxicity testing, low efficacy and lack of transparency may endanger the lives of iron loaded patients. *Hemoglobin* 35:301–311.
- Lewis RE, et al. 2011. Activity of deferasirox in *Mucorales*: influences of species and exogenous iron. *Antimicrob. Agents Chemother.* 55:411–413.
- Loyevsky M, et al. 1999. Chelation of iron within the erythrocytic *Plasmodium falciparum* parasite by iron chelators. *Mol. Biochem. Parasitol.* 101:43–59.
- Neupane GP, Kim DM. 2010. *In vitro* time-kill activities of ciprofloxacin alone and in combination with the iron chelator deferasirox against *Vibrio vulnificus*. *Eur. J. Clin. Microbiol. Infect. Dis.* 29:407–410.
- Nunes A, et al. 2010. Fluorescent 3-hydroxy-4-pyridinone hexadentate iron chelators: intracellular distribution and the relevance to antimicrobial properties. *J. Biol. Inorg. Chem.* 15:861–877.
- Qiu DH, Huang ZL, Zhou T, Shen C, Hider RC. 2011. *In vitro* inhibition of bacterial growth by iron chelators. *FEMS Microbiol. Lett.* 314:107–111.
- Reyes HM, Tingle EJ, Fennes AZ, Spiegel J, Burton EC. 2008. Pulmonary invasive mucormycosis in a patient with secondary iron overload following deferoxamine therapy. *Proc. (Bayl. Univ. Med. Cent.)* 21:378–381.
- Sandiumenge A, Rello J. 2012. Ventilator-associated pneumonia caused by ESKAPE organisms: cause, clinical features, and management. *Curr. Opin. Pulm. Med.* 18:187–193.
- Shachar DB, Kahana N, Kampel V, Warshawsky A, Youdim MB. 2004. Neuroprotection by a novel brain permeable iron chelator, VK-28, against 6-hydroxydopamine lesion in rats. *Neuropharmacology* 46:254–263.
- Soummer A, et al. 2008. Failure of deferasirox, an iron chelator agent, combined with antifungals in a case of severe zygomycosis. *Antimicrob. Agents Chemother.* 52:1585–1586.
- Spellberg B, et al. 2008. The epidemic of antibiotic-resistant infections: a call to action for the medical community from the Infectious Diseases Society of America. *Clin. Infect. Dis.* 46:155–164.
- Tam TF, et al. February 2011. Process for the manufacture of 3-hydroxy-n-alkyl-1-cycloalkyl-6-alkyl-4-oxo-1,4-dihydropyridine-2-carboxamide and its related analogues. Apotex, Inc. US patent 7,893,269.
- Tribble DR, et al. 2011. Infection-associated clinical outcomes in hospitalized medical evacuees after traumatic injury: trauma infectious disease outcome study. *J. Trauma* 71:S33–S42.
- van Asbeck BS, et al. 2001. Anti-HIV effect of iron chelators: different mechanisms involved. *J. Clin. Virol.* 20:141–147.
- Walcourt A, Loyevsky M, Lovejoy DB, Gordeuk VR, Richardson DR. 2004. Novel aroylhydrazones and thiosemicarbazones iron chelators with anti-malarial activity against chloroquine-resistant and -sensitive parasites. *Int. J. Biochem. Cell Biol.* 36:401–407.
- White RL, Burgess DS, Manduru M, Bosso JA. 1996. Comparison of three different *in vitro* methods of detecting synergy: time-kill, checkerboard, and E test. *Antimicrob. Agents Chemother.* 40:1914–1918.
- Zarembek KA, Cruz AR, Huang CY, Gallin JI. 2009. Antifungal activities of natural and synthetic iron chelators alone and in combination with azole and polyene antibiotics against *Aspergillus fumigatus*. *Antimicrob. Agents Chemother.* 53:2654–2656.
- Zheng H, Gal S, Weiner LM, Bar-Am Warshawsky OA, Fridkin M, Youdim MB. 2005. Novel multifunctional neuroprotective iron chelator-monoamine oxidase inhibitor drugs for neurodegenerative diseases: *in vitro* studies on antioxidant activity, prevention of lipid peroxide formation and monoamine oxidase inhibition. *J. Neurochem.* 95:68–78.